

## <u>Lipodissolve Mesotherapy Consent Form</u>

l,	, voluntarily consent to undergo Mesotherapy
ė –	ers or other certified body contourist, nurses,
physician associates or qualified sta	ff members employed by the practice.
I understand that Mesotherapy can b treatment for the following; Reducti	oe used for many reasons and I want to have on of localized fat of
one (1) treatment is required. Lunde	rtreatment of which I understand that more than restand that the treatment requires many small reated. I understand that the administration of med needed.
I understand that there are some risk possible risks with Mesotherapy:	as with any procedure. The following is the list of
<ul> <li>Itching and burning lastin</li> <li>Scarring of the skin is unl</li> <li>Nausea, dizziness and pos occur.</li> </ul>	ne injections. e injection site. of the skin (transient or permanent) ng 20 minutes to a few hours
	nmended to reduce bruising, swelling and um of 3 days prior to scheduled treatment.
medications and give consent to its u medicine is not an exact science; the results of my treatments. I understa	at I have been informed about the above sein my treatment. I know that the practice of erefore, no guarantee can be made as to the nd that this treatment is strictly for cosmetic insurance. I understand that I am responsible service.
By my signature, I certify that I have t this form and the disclosures listed	choroughly read and understand the contents of above were made to me.
Patient's Signature	Date



## **Mesotherapy Post-Treatment Instructions**

- Immediately after the treatment, the most commonly reported side-effects were temporary redness, bruising and swelling at the injection site. These effects typically resolve within 5 to 10 days. Cold or ice compresses may be used immediately after treatment to reduce swelling.
- Continue taking Arnica Montana up to 7 days after each treatment to decrease bruising and inflammation.
- Apply 1% Hydrocortisone cream or Benadryl spray or gel on treated areas to reduce itching or redness.
- To minimize bruising, avoid Aspirin, Anti-inflammatory drugs, Gingko biloba, Garlic, Flaxseed Oil, Vitamin E, Alcohol, spicy food, salty food and cigarettes 48 hours to 1 week after your treatment.
- It is normal to feel "firmness" in the injection site first day after treatment. In some cases, a lumpy formation can be felt on the injected area. If necessary, massage area gently 2-3 times a day up to 72 hours.
- Do not exercise for 24 hours after treatment. Avoid strenuous exercises, sunbathing or tanning.
- Apply sun block and protect skin from sunlight.
- For treatment of neck areas, sleep with head elevated (3-4 pillows), and wear some compression under chin (scarf or head band).
- Call us immediately if you start experiencing these symptoms or develop any persistent side effect at .

Patient's Signature	Date	

R. 5/2016